



Syncope Unit: rationale and requirement – the European Heart Rhythm Association position statement endorsed by the Heart Rhythm Society

Rose Anne Kenny (Chairperson, Ireland), Michele Brignole (Co-chairperson, Italy), Gheorghe-Andrei Dan (Romania), Jean Claude Deharo (France), J. Gert van Dijk (The Netherlands), Colin Doherty (Ireland), Mohamed Hamdan (USA), Angel Moya (Spain), Steve W. Parry (UK), Richard Sutton (UK), Andrea Ungar (Italy), and Wouter Wieling (The Netherlands)

External contributors to the Task Force: Mehran Asgari (UK), Gonzalo Baron-Esquivias (Spain), Jean-Jacques Blanc (France), Ivo Casagrande (Italy), Conal Cunningham (Ireland), Artur Fedorowski (Sweden), Raffello Furlan (Italy), Nicholas Gall (UK), Frederik J. De Lange (The Netherlands), Geraldine McMahon (Ireland), Peter Mitro (Slovakia), Artur Pietrucha (Poland), Cristian Podoleanu (Romania), Antonio Raviele (Italy), David Benditt (USA), Andrew Krahn (Canada), Carlos Arturo Morillo (Canada), Brian Olshansky (USA), Satish Raj (USA), Robert Sheldon (Canada), Win Kuang Shen (USA), Benjamin Sun (USA), Denise Hachul (Brazil), Haruhiko Abe (Japan), and Toshiyuki Furukawa (Japan)

Document reviewers: Review coordinators, Bulent Gorenek (Turkey), Gregory Y. H. Lip (UK), Michael Glikson (Israel), Philippe Ritter (France), Jodie Hurwitz (USA), Robert Macfadyen (Australia), Andrew Rankin (UK), Luis Mont (Spain), Jesper Svendsen (Denmark), Fred Kusumoto (USA), Mitchell Cohen (USA), and Irene Savelieva (on behalf of EP-Europace, UK)

Keywords

Syncope • Neurocardiogenic syncope • Transient loss of consciousness • Faint • Fall • Seizure • Syncope Unit

Table of Contents

Scope of the document	2	Existing models for syncope management.	5
Definitions	2	Current situation	7
Definition of syncope and transient loss of consciousness	2	General attributes of a Syncope Unit.	7
Definition of a Syncope Unit	2	Structure of the European Heart Rhythm Association	
Rationale for a Syncope Unit.	2	Syncope Unit.	8
Expected benefit and barriers to setting-up a Syncope Unit	2	Competence.	9
Syncope Unit reduces underdiagnosis and misdiagnosis		Considerations.	9
of syncope	2	Syncope specialist.	9
Syncope Unit reduces hospitalization	3	Notes on training	10
Syncope Unit reduces cost of syncope.	3	Comprehensive care.	10
Structure of European Heart Rhythm Association Syncope Unit	5	Referrals, access criteria	10

Management modalities	11
Final report and follow-up	12
Quality indicators	13
References	15
Appendix	16

Scope of the document

The European Society of Cardiology (ESC) has played an important role in advancing our understanding of the causes, optimal investigation, and management of syncope through publication of practice guidelines in 2001, 2004, and 2009.^{1–3} The 2009 ESC guidelines recommend the establishment of formal Syncope Units (SUs)—either virtual or physical site within a hospital or clinic facility—with access to syncope specialists and specialized equipment.³ In response, this position statement by the European Heart Rhythm Association (EHRA) endorsed by the Heart Rhythm Society (HRS) offers a pragmatic approach to the *rationale and requirement for an SU*, based on specialist consensus, existing practice and scientific evidence (see Appendix).

The panel consists of specialists who have experience in developing and leading such units representing cardiology, geriatric and general internal medicine, neurology, and emergency medicine.

This document is addressed to physicians and others in administration, who are interested in establishing an SU in their hospital, so that they can meet the standards proposed by ESC-EHRA-HRS.^{1–3}

Definitions

Definition of syncope and transient loss of consciousness

Syncope is a transient loss of consciousness (T-LOC) due to transient global cerebral hypoperfusion, and is characterized by rapid onset, short duration, and spontaneous complete recovery. This definition of syncope has been developed by the Task Force for the Diagnosis and Management of Syncope of the ESC and endorsed by the EHRA, European Heart Failure Association and European Heart Rhythm Society. Transient loss of consciousness is a term that encompasses all disorders characterized by self-limited LOC, irrespective of mechanism.^{1–3} By including the mechanism of unconsciousness, i.e. transient global cerebral hypoperfusion, the current syncope definition excludes other causes of T-LOC such as epileptic seizures and concussion, as well as certain common syncope mimics, such as psychogenic pseudosyncope.

Definition of a Syncope Unit

An SU is a facility featuring a standardized approach to the diagnosis and management of T-LOC and related symptoms, with dedicated staff and access to appropriate diagnostics and therapies. The SU should also take the lead in educating and training clinicians who encounter syncope. Even if the most appropriate term describing such an organization should be the more general T-LOC Unit (or

Faint Unit), this Task Force decided to maintain the term of SU, because it is most frequently used worldwide. This Position Paper is a pragmatic approach to outline the constituents of an SU and assist target groups with the current available necessary information. The authors emphasize that there is, at present, insufficient available evidence whether an SU (examples of a number of models are detailed later in the document) is superior in efficiency or outcomes to a syncope specialist⁴ or newer technologically driven models of syncope management.⁵ We anticipate that the Position Paper will stimulate structured research to determine best practice models for T-LOC evaluation in different settings and cultures.

Rationale for a Syncope Unit

Expected benefit and barriers to setting-up a Syncope Unit

Syncope is a common medical problem that can be debilitating and associated with high healthcare costs.^{6–9} There is wide variation in practice of syncope evaluation, and wide variation in adoption of recommendations from published guidelines.^{10,11} The absence of a systematic approach to T-LOC incurs higher health and social care costs, unnecessary hospitalizations, and diagnostic procedures, prolongation of hospital stays, lower diagnostic rates, and higher rates of symptom recurrences. Therefore, a systematic approach, by a dedicated service (an SU), equipped to evaluate and manage this common problem may ensure better management of T-LOC, from risk stratification to diagnosis, therapy and follow-up (Table 1).

Despite the recommendation from the ESC,^{2,3} SUs are not widely established in clinical practice. Possible reasons for this are outlined in Table 2. Barriers to establishing an SU include lack of resources, lack of trained dedicated staff, and complex presentations to multiple settings, necessitating involvement from multiple disciplines. When developing a case of need for the SU, individual practices may not be able to access detailed information to inform fully the economic cost and resource requirements necessary and this can make the justification for realignment of resources challenging. This document will assist practitioners to develop a model best suited to local requirements.

Syncope Unit reduces underdiagnosis and misdiagnosis of syncope

An Internet search of the phrase 'Misdiagnosis of Syncope', reveals about equal numbers of search hits from three perspectives: those who approach the problem from the perspective of the over diagnosis of epilepsy,²⁸ the underdiagnosis of syncope,²⁹ or legal firms soliciting business from victims of either. When it comes to the underdiagnosis and misdiagnosis of syncope, are estimated to be as high as 40%.^{17,28–32} Underdiagnosis and misdiagnosis have been reported in both outpatient and emergency settings. Although there are no large-scale randomized trials comparing misdiagnosis of syncope in SUs to usual care, smaller cohort studies confirm

Table 1 Expected benefits of Syncope Unit

Expected benefits
Specialist opinion for patients
Early accurate and efficient diagnosis
Timely treatment
Better application of recommended guidelines
Less duplication and fragmentation of services
Single source of communication for all stakeholders
Shorter length of stay for hospital inpatients
Reduction of total care costs
Better systems for monitoring and evaluation of practice at local, national, and international level
Better quality control at local, national and international level
Access to harmonized data across different hospitals
High quality, evidence-based data for research
Evidence-based innovation in diagnosis, treatments and healthcare model

Table 2 Barriers to establishing the Syncope Unit

Barriers to establishing an SU
Lack of awareness of the benefits of an SU due to inadequate research trials comparing SUs to normal practice
Underestimation of consequences of syncope
Lack of awareness of benefit of an SU on quality-of-life
Low numbers of syncope specialists
Lack of formal syncope training programmes
Wide age range from paediatric to oldest patients
Skill sets required in a number of domains such as cardiology, geriatrics, paediatrics, physiology, neurology, and psychiatry
Syncope not a recognized subspecialty
Reluctance to introduce innovative proposals
Necessity to engage multiple stakeholders
Inadequate reimbursement of syncope core management
New economic cost models required to evaluate an SU
Fear of increasing costs by the development of a new structure instead of reducing them

high rates of misdiagnosis with usual care and the benefits of a structured approach to diagnosis.^{9,17,20,33}

Syncope Unit reduces hospitalization

In hospitals without an SU, T-LOC evaluation and management is more frequently carried out as an inpatient rather than an outpatient service. In other words, patients are preferentially admitted to emergency services rather than evaluated and managed as outpatients (Table 3). In one study, the average length of stay for acute admissions due to syncope or collapse was two-fold higher in hospitals without an SU.^{12,15,34,35}

The Syncope Evaluation in the Emergency Department Study (SEEDS)¹⁵ randomized intermediate-risk patients to an Emergency Department (ED)-based explicit syncope protocol vs. routine inpatient admission. Hospital admissions were reduced by 56%, and total patient-hospital days were reduced by 54%. In the Emergency Department Observation Syncope Protocol (EDOSP) trial,²⁶ patients randomized to an ED observation protocol experienced a 77% reduction in hospitalization and a 40% reduction in hospital length of stay.²⁶ An integrated model of Short Observation stay in ED, coupled with fast track to an SU allowed a reduction of the admission rate to 29% with 20% of patients being discharged after a short observation in ED, 20% fast-tracked to the SU and 31% directly discharged.³⁶

Syncope Unit reduces cost of syncope

Cost estimates

Several studies investigated costs of syncope as they appear in national or hospital records (Table 3).^{24,34,37,38} In the USA, in 2004, the mean cost for a syncope-related admission was \$5400 (95% CI: \$5100–5600) with a total annual cost of \$2.4 billion,³² similar to asthma (\$2.8 billion) and HIV (\$2.2 billion). In Italy, in 2006,¹⁷ the mean cost for a syncope-related admission was €2785 ± 2168; hospital costs accounted for about three quarters of total costs. The cost

per patient discharged from ED was €180 ± 63. In Spain, in 2006,⁹ the overall cost (which included stay, diagnosis, and treatment) per admitted patient was €11 158 (range: 1651–31 762). In the outpatient setting, the cost is similarly alarmingly high due to significant variability in practice and the use of unnecessary tests.^{6,19,39} In one study,⁴⁰ patients with unexplained syncope had a median of 13 non-diagnostic tests performed (range: 9–20) before receiving an implantable loop recorder.

Cost reductions

Various hospitals organized syncope care through the creation of SUs, where a solid conceptual framework with clearly delineated diagnostic procedures is implemented. The primary outcome was an increase in the rate of diagnostic yield,^{1–3} and a reduction in costs primarily by reducing the number of admissions, duration of hospital stay, and the number of unnecessary tests,^{12,13,15,17,19,21–23,25,41,42} with few exceptions.¹⁴ Indeed, Brignole *et al.*¹⁷ have shown a 19% reduction in cost per patient and a 29% reduction in cost per diagnosis in the standardized care group when compared with the conventional approach. The EDOSP randomized trial²⁶ included an explicit cost analysis. Hospital facility costs were \$629 less in the observation unit group compared with the routine admission group. Rates of diagnostic testing and specialty consultation were similar in both groups; therefore, cost savings were due to reduction in hospital length-of-stay.

Expected benefit

Establishing an SU should benefit three parties: (i) patients by increasing the rate of correct diagnosis, (ii) health payers by reducing total cost per patient and diagnosis, and (iii) hospitals by increasing their market share via the added value proposition. In healthcare systems based on payment per test or per medical action, an SU may reduce income through a reduction of tests. The expected benefit should focus on improved healthcare.

Table 3 Comparison between systematic evaluation and conventional management of syncope in controlled studies

Source	Intervention	Comparison setting	Number of patients	Main results (experimental vs. conventional group)
Kenny <i>et al.</i> ¹² UK	Day-case falls and syncope unit in adults	Bed day activity in hospitals with (E) vs. hospitals without day-care facility (C)	–	64% reduction in admissions in hospital with falls and syncope
Brignole <i>et al.</i> ¹³ Italy	In-hospital Syncope Unit within the Department of Cardiology	6 hospitals with Syncope Unit (E) vs. 6 hospitals without (C) Syncope Unit	274 (C) vs. 279 (E)	11% reduction in unexplained syncope 8% reduction of tests 12% reduction in admissions
Farwell and Sulke ¹⁴ UK	Diagnostic/management protocol for syncope	One year with protocol (E) vs. previous year in one hospital	660 (C) vs. 421 (E)	166% increase in costs per patient: despite improved diagnosis, inappropriate investigation and admission still occurred
Shen <i>et al.</i> ¹⁵ USA	Syncope Unit in the ED	Rate of admission and diagnosis in patients randomized to Syncope Units (E) vs. standard of care (C)	51 (C) vs. 52 (E)	63% reduction in unexplained syncope 56% reduction in admissions
Blanc <i>et al.</i> ¹⁶ France	Education of ED physicians on T-LOC/ syncope	One year before (C) vs. 1 year after education (E)	454 (C) vs. 524 (E)	11% reduction in unexplained syncope –25% reduction in admissions
Brignole <i>et al.</i> ¹⁷ Italy	Standardized care using ESC guidelines	28 hospitals with standard care (C) vs. 19 with standardized care (E)	929 (C) vs. 745 (E)	25% reduction in unexplained syncope 24% reduction of tests 17% reduction in admissions 19% reduction of costs per patient
Parry <i>et al.</i> ¹⁸ UK	Education through management algorithm for acute medical services; effect on patient admitted for falls and syncope	One-month period before (C) vs. 1-month period a year later (E)	41 (C) vs. 31 (E)	2% reduction in unexplained syncope
Ammirati <i>et al.</i> ¹⁹ Italy	Syncope Unit	Unexplained syncope referred to Syncope Unit; period before patients visited Syncope Unit (C) vs. after visit (E)	96	82% reduction in unexplained syncope –85% reduction of costs per patient
Fedorowski <i>et al.</i> ²⁰ Sweden	Syncope Unit	Unexplained syncope patients discharged from ED or hospital ward (C) vs. the same patients evaluated by SU (E)	101	87% reduction in unexplained syncope
McCarthy <i>et al.</i> ²¹ Ireland	Using ESC Guidelines	Utilization of resources in ED (C) vs. re-evaluation of same patients using ESC guidelines; 6-month period	214	54% reduction in admissions
Daccarett <i>et al.</i> ²² USA	ESC Guidelines incorporated in 'Faint-Algorithm'	Retrospective assessment of ED admissions	254	52% reduction in admissions
Shin <i>et al.</i> ²³ South Korea	Standardized ED protocol for syncope based on ESC guidelines	Period before (C) and after (E) standardization	116 (C) vs. 128 (E)	28% reduction in unexplained syncope 39% reduction in admissions 32% reduction of costs per patient
Sun ²⁴ USA	Up to 24 h observation ED	ED syncope presentation with usual care (C) or observation period (E) for intermediate-risk patients	62 (C) vs. 62 (E)	84% reduction in admissions 42% reduction of costs per patient
Sanders <i>et al.</i> ²⁵ USA	Standardized care implemented in Faint and Fall Clinic vs. historical control	Standardized care (E) vs. historical control (C)	100 (C) vs. 154 (E)	22% reduction in unexplained syncope 80% reduction in admissions
Sun <i>et al.</i> ²⁶ USA	Observation Unit in five EDs	Rate of admission and costs in patients >50 years, randomized to observation unit (E) vs. standard of care (C)	62 (C) vs. 62 (E)	84% reduction in admissions 629 \$ reduction in index hospital costs
Raucci <i>et al.</i> ²⁷ Italy	Standardized care implemented of paediatric guidelines vs. historical control in ED	Two years with protocol (E) vs. previous 2 years in one hospital	470 (C) vs. 603 (E)	72% reduction in unexplained syncope 54% reduction in admissions

E, experimental group; C, control group; ED, Emergency Department.

There may be cost savings, but these will be system dependent and thus vary. The cost benefit of a syncope specialist or an SU in different settings and different healthcare systems has not been exposed to rigorous economic and scientific scrutiny. Further research is required to determine resource outcomes and the authors acknowledge the limitations of the current knowledge base and recognize that service models may be influenced by local circumstances.

Structure of European Heart Rhythm Association Syncope Unit

Existing models for syncope management

Syncope management organization may differ widely among healthcare systems and from hospital to hospital. A review of published data on organization and impact on outcomes of models of care may guide SU design and implementation for a given environment. *Table 4* summarizes these data. We acknowledge that overlap exists between these models.

A functional Syncope Unit located in a cardiology department^{13,17,20,25,39,43–45}

In this model, introduced in Italy and adopted in other countries such as Sweden, Portugal, USA, and France, the SU is supervised by cardiologists, supported by dedicated personnel with expertise in syncope. Patients access the SU mainly as outpatient or via the emergency room. This model has evolved into a 'virtual' unit based on the expertise of a limited team and in some instances on web-based decision-making software.⁵² Access to a specialist is regarded as essential. A specialist can be accessed by any means, e.g. telephone. The Evaluation of Guidelines in SYncope Study-2 (EGSYS-2)¹⁷ showed a sharp decrease in the overall cost of care driven by a reduction in average cost per patient of 19% and average cost per diagnosis of 29%. Seventy-one Italian hospitals now have an SU that has been certified by peer-review members of GIMSI (Gruppo Italiano Multidisciplinare per lo studio della Sincope, www.gimsi.it). Similar models have been described in other departments, e.g. geriatric and internal medicine; the organization is basically the same with formalized fast-tracking processes to cardiological testing. In a few instances, the SU includes also a short observation stay as part of an internal protocol for risk stratification of intermediate risk ED patients.³⁶

The Day-Care Syncope Evaluation Unit and the Falls and Syncope Services

This model was first developed in Newcastle, UK¹² and takes the form of an outpatient, day-care facility located in a general hospital. The service provides a multidisciplinary approach based on the application of evidence-based diagnostic algorithms to patients with falls (for older patients) and T-LOC of suspected syncopal nature (all adult age groups). Close liaison exists with acute medical in-patient and ED services. After a consultation with an emergency physician, geriatrician, internist, or general practitioner, patients have access to non-invasive diagnostic testing, occupational and physiotherapy,

and supplementary examinations supported by specialist nurses. There is close cooperation and consultation with cardiology and neurology services for further diagnostics and treatments. In line with this experience, Falls and Syncope Services for older people have evolved by defining protocols and educational methods for inpatients³⁵ and outpatients.⁴⁶

The Rapid Access Blackouts Clinic

This type of SU functions as a 'referral centre' for patients with T-LOC.^{47,48} As such, it is positioned between first response and specialist referrals. It is led jointly by a cardiologist and a neurologist. The aim is to provide rapid access to clinical and ECG assessment in order to screen patients. The SU is run by nurses specialized in epilepsy, cardiac arrhythmias, or geriatrics. Patients referred by general practitioners or emergency services complete a triage questionnaire: 60 standard questions/data detailing characteristics of falls, syncope, and epilepsy, which is analysed by the nurses. A cardiologist may be consulted for interpretation of tests, notably the ECG. Following this evaluation, patients are referred, as appropriate, to a cardiologist, neurologist, geriatrician, general practitioner, or psychologist. Continuity of care is ensured by maintaining and sharing a database for all stakeholders.

Tertiary referral Syncope Unit^{49,53}

This model is centred on one syncope specialist, a neurologist, internist, or cardiologist, who mostly sees tertiary T-LOC referrals from neurologists and cardiologists. The SU consists of an outpatient clinic and a core laboratory performing tilt table test, carotid sinus massage, cardiovascular autonomic tests, and ambulatory BP monitoring. The tertiary character means that other ancillary tests have already been performed, and that the case mix concerns low rates of cardiac syncope and of unexplained syncope, but high rates of reflex syncope, psychogenic pseudosyncope, and epilepsy (see 'Competence' section).

The Syncope Observational Unit in the Emergency Department

This type of SU is described in the SEEDS study.¹⁵ This single-centre prospective randomized study evaluated a standardized unit incorporated in the ED of a university tertiary care hospital, compared with conventional care, in a group of syncope patients considered to be at 'intermediate' cardiovascular risk. After 6 h of monitoring and tests in the ED which included regular orthostatic blood pressure measurements, tilt table test and carotid sinus massage upon physician's request and electrophysiologist's consultation when requested, patients without indication for hospitalization were offered rapid outpatient follow-up consultation. This study showed, in 51 consecutive patients randomized to SU evaluation, an improvement in diagnostic yield compared with conventional care (67 vs. 10%) and a decrease in hospital admission (43 vs. 98%), but no changes in the average length of stay. The model did not reduce 2-year mortality nor syncopal recurrences.

Using a protocol with an initial evaluation similar to that described in the ESC Guidelines and an observational unit, for up to 24 h, within the ED, a Spanish group has achieved a diagnosis in 78% of patients

Table 4 Various existing models for syncope management from published data of comparison between systematic evaluation and conventional management in controlled studies

	The functional Syncope Unit in a cardiology department	The Day-Care Syncope Evaluation Unit and Fall and Syncope Services	The Rapid Access Blackouts Triage Clinic (T-LOC Triage Clinic)	Tertiary referral SU	The Syncope Observational Unit in the ED	The web-based standardized care pathway for Faint and Fall patients (Faint and Fall Clinic)
References	13,17,20,25,36,39,41,43–45	12,35,46	47,48	49,50	15,26,51	5,25
Location	Cardiology department/ outpatient clinic	Outpatient clinic	Outpatient clinic	Outpatient clinic	ED	Outpatient clinic
Management	Cardiologist with rapid access to other specialists	Geriatrician/internist	Specialized nurses (arrhythmias, falls, epilepsy) Supervision: cardiologist/ neurologist	One syncope specialist (neurologist, internist, cardiologist)	Experienced emergency physician	Cardiologist and geriatrician with rapid access to a neurologist
Support	Trained nurses	Other specialists and general practitioners, specialized nurses	Cardiologists	Technicians, specialized nurses	Specialized nurses, electrophysiologist's, other specialists	Nurse practitioner
Referral	Outpatients, fast track from ED, other departments	Community, ED, other departments	General practitioners, specialists (cardiology, neurology), ED	Most referrals from cardiologists and neurologists	ED (only intermediate risk patients were included in the SEEDS and EDOSP studies)	Outpatients and ED
Organization	Functional unit in the hospital	Day-care multidisciplinary medical approach, specialized nurses	Rapid assessment outpatient clinic	Outpatient clinic	6–24 h of observation	Fixed unit with rapid access
Tools	Guidelines-based flowcharts, software	Specialist visits, non-invasive tests, occupational activities	Web-based questionnaire	History taking, reappraisal of the case	ECG and BP monitoring, non-invasive tests and electrophysiologist's consultations. Rapid FU appointment if discharged	Web-based decision-making software
Core laboratory tests	CSM, TT with beat-to-beat measurement, ILR and ELR, ambulatory BP monitoring	CSM, TT with beat-to-beat measurement, ILR, and ELR	History, physical examination, ECG, ILR	TT, CSM, autonomic tests, ambulatory BP and ECG monitoring, ILR	Laboratory tests, CSM, TT	Cardiac imaging, stress tests, TT, CSM, electrophysiological study, Holter, ELR, and ILR
Impact on outcomes: methodology	Lowering of hospital admissions and costs, improvement of the diagnostic yield	Lowering of costs driven by hospital admissions and readmissions, improvement of the diagnostic yield and access to tests	Rapid diagnosis and triage, lowering of readmissions for T-LOCs	Compared with other SUs low rates of unexplained syncope and of cardiac syncope, high rates of psychogenic pseudosyncope and complex reflex syncope	Higher and earlier number of suspected diagnosis, lower hospital admissions and patient-hospital days	Decrease in hospital admissions, higher rate of diagnosis at 45 days, less utilization of costly tests and consultations

BP, blood pressure; CSM, carotid sinus massage; TT, tilt table test; ILR, implantable loop recorder; ELR, external loop recorder; T-LOC, transient loss of consciousness; ED, emergency department; ECG, electrocardiogram.

presenting as emergencies with only 10% of syncope patients being admitted to hospital in this model.⁵¹

Recently, the EDOSP study²⁶ evaluated an ED observation protocol at five sites, including university, community, and public hospitals. Patients at 'intermediate' risk were randomized either to an explicit ED-based observational unit protocol vs. routine care. The observational unit protocol included up to 24 h of cardiac monitoring and echocardiogram for selected patients. In 124 randomized patients, there were reductions in hospitalizations rate (15 vs. 92%), length-of-stay (29 vs. 47 h), and hospital costs were \$629 lower than the admission group. There were no differences in safety events (i.e. serious 30-day outcomes occurring after hospital discharge), quality-of-life, or costs. The EDOSP study generalizes the SEEDS findings to a diverse set of hospitals, and includes novel assessments of patient-centred outcomes and a formal economic analysis.

The Faint and Fall Clinic

The Faint and Fall Clinic offers a multidisciplinary approach to patients presenting with fainting spells or falls. Patients are evaluated by advanced nurse practitioners and then seen by a cardiologist or a geriatrician with rapid access to a neurologist, as needed. Providers in the clinic use a web-based interactive software that integrates the most recent guidelines for risk assessment and management of patients with T-LOC.^{5,25}

Current situation

The UK has a growing number of SUs, which are listed by STARS (Syncope Trust and Anoxic Reflex Seizures), a charitable organization providing information to patients about syncope and related conditions, www.stars.org. Italy has a growing number of SUs which are listed by GIMSI (Gruppo Italiano Multidisciplinare per lo studio della Sincope), www.gimsi.it. These information sites include available SUs that are geographically close to the enquiring patient. In Ireland, USA, Canada, Spain, Portugal, France, Netherlands, Sweden, Japan, Brazil, Romania, Poland, Slovakia, and other countries there are similar developments.

In summary, whatever the SU model, the key elements are rapid access to syncope expertise in trained, dedicated staff, together with the utilization of standardized algorithms. The European Heart Rhythm Association considers that SUs should be widely available in Europe. Their aims and structure should be in line with one or other of the models reported permits each hospital to develop their own model to suit its particular environment.

General attributes of a Syncope Unit

The SU can be virtual or based on a pre-defined location such as a unit associated with the ED, an ambulatory clinic, or employs a combination of approaches. The model of SU should be the best fit for local practice. Because of the extensive differential diagnoses and high prevalence of syncope in older patients, the skill mix of SU staff should include some training/knowledge of common disorders that cause or mimic syncope: commonly cardiovascular, neurologic, geriatric, and psychiatric disorders.

As no single syncope care service model is suitable for all health-care systems, the following is a list of some of the *important features* to consider when establishing an SU:

- Structure of SU
 - The model of care delivery should be appropriate to local resources and local specialities while ensuring implementation of published practice guidelines.
 - Models of care delivery vary from a single one-site-one-stop syncope facility to a wider multifaceted practice in which several specialists are involved in syncope management.
 - The SU can be a single site facility or virtual model with mobile team.
- Stakeholders
 - All key stakeholders should be involved in the earliest stages of development and implementation of the SU.
- It is essential to establish a mechanism through which regular communication can be established with all stakeholders (i.e. patients, referring physicians, hospital/clinic management, consultant physicians, nurses, and other allied medical professionals) in order to ensure an ongoing consensus for and understanding of proposed management strategies. This mechanism includes the implications of and implementation of published guidelines. Stakeholders may be staff from the ED, neurology, general internal medicine, orthopaedic surgery, geriatric medicine, psychiatry, and ear, nose, and throat (ENT), paediatric departments in addition to cardiology.
- This should also include agreed measurement metrics of performance in order that early recognition of variation is achieved; thereby re-calibration of the model can take place in a timely fashion. These measures should include outcome measures, process measures, and balancing measures.
- A clear diagnostic and therapeutic pathway provides a framework, which is fundamental to enable new evidence to be incorporated into the model seamlessly.
- Management
 - The management strategy should be agreed on and practiced by all practitioners (encompassing a range of specialities) involved in syncope management.
- Patient case mix
 - The age range and symptom characteristics of patients appropriate for syncope investigation should be determined in advance. Some facilities are prepared to evaluate both paediatric and adult patients, whereas others limit practice to adult or paediatric cases. A wide age range is encouraged.
- Referral sources
 - Potential referral sources should be taken into consideration. Referral can be directly from family practitioners, from the ED, from occupational physicians, from hospital admissions, and from patients in institutional settings. The scope of referral source has implications for resources and skill mix.
- SU—skill mix and staffing
 - There are existing models in which cardiologists (commonly with an interest in cardiac pacing and electrophysiology),

- neurologists (commonly with an interest in autonomic disorders and/or epilepsy), internists (commonly with an interest in cardiovascular physiology and autonomic disorders), emergency doctors, and geriatricians (commonly with an interest in age-related cardiology or falls) each may lead syncope facilities. There is no evidence for superiority of any model.
- The skill mix (i.e. the types of professional/specialities) required to staff the facility depends on the extent to which screening of referrals occurs before presentation at the facility. For example, if referrals are directly from the community a broader skill mix than cardiology is required. Under these circumstances, other disorders such as epilepsy, autonomic disorders, neurodegenerative disorders, metabolic disorders, and falls are also common.
 - When establishing a unit the lead clinician should have knowledge of the catchment area for referrals and projected volume in order to estimate staff requirements and to tailor the scope of referrals to available resources. The volume of activity and the number of personnel largely vary based on the model of SU and local organization. No empirical figures can therefore be given. However, since the aim of this document is to provide practical advice to stakeholders who are interested in setting-up an SU, as a general guide, this Task Force believes

that the following figures should be of help. In the Italian experience,⁴¹ 163 patients per 100 000 inhabitants per year were referred to the local SU. They performed an average of 2.9 tests per patient. Patients will be followed-up from a minimum of one to multiple visits. Multiple visits (on-site or by means of telemedicine) were necessary especially in the case of patients with unexplained syncope undergoing prolonged monitoring or patients who had received device therapies (pacemaker clinic, etc.). Thus, this Task Force estimates that one syncope specialist and one technician need to work the equivalent of one full working day per week for every 100 000 inhabitants of the catchment area.

- SU staff should provide ongoing education and training in syncope diagnosis, investigation and management to primary and secondary care colleagues who deal with this symptom in their day-to-day practice.

Structure of the European Heart Rhythm Association Syncope Unit

The proposed structure of the EHRA SU is shown in the *Consensus Statement 1*. The role of physician and staff in performing procedures and tests is shown in *Table 5*.

Table 5 The role of physician and staff in performing procedures and tests

Procedure or test	SU physician	SU staff	Non-SU personnel
History taking	x		
Structured history taking (e.g. application of software technologies and algorithms)		x	
12-Lead ECG		x	
Blood tests		x	
Echocardiogram and imaging			x
Carotid sinus massage	x		
Active standing test		x	
Tilt table test	x ^a	x	
Basic autonomic function test	x	x	
ECG monitoring (Holter, external loop recorder): administration and interpretation	x	x	
Implantable loop recorder	x	x ^b	
Remote monitoring		x	
Other cardiac tests (stress test, electrophysiological study, angiograms)			x
Neurological tests (CT, MRI, EEG, video-EEG)			x
Pacemaker and ICD implantation, catheter ablation			x
Patient's education, biofeedback training ^c , and instruction sheet on counter pressure manoeuvres	x	x	
Final report and clinic note	x		
Communication with patients, referring physicians and stakeholders	x	x	
Follow-up	x	x	

CT, computed tomography; ECG, electrocardiogram; EEG, electroencephalogram; ICD, implantable cardioverter-defibrillator; MRI, magnetic resonance imaging; SU, Syncope Unit.

^aPhysician need not be in the room, but a physician adequately trained in resuscitation needs to be in the area of the test.

^bCurrent practice limited to few countries.

^cBiofeedback means that the training session of the counter pressure manoeuvres consists of biofeedback training using a continuous blood pressure monitor. Each manoeuvre is demonstrated and explained. The manoeuvres are practiced under supervision, with immediate feedback of the recordings to gain optimal performance.⁵⁴

Consensus Statement 1—Structure of the EHRA SU

Staffing of an SU is composed of:

- (1) One or more physicians of any specialty who are syncope specialists. Owing to the multidisciplinary nature of T-LOC management, each SU should identify specific specialists for SU and for consultancies.
- (2) A team comprising professionals who will advance the care of syncope patients. These may be physicians, specialized/trained nurses, or others who bring multidisciplinary skills to the facility, coupled with administrative support. The roles played by members of the team may vary according to local circumstances and individual skill. Nurses may be expected to take very important roles including initial evaluation of patients, follow-up clinic assessments, selection of investigations including tilt testing and implantation/insertion of ECG loop recorders according to pre-defined protocols and local regulations.
- (3) Given that the SU is integrated within a hospital organization, syncope specialists, and staff are not necessarily employed full-time, but frequently have other duties depending on the volume of activity in the SU.

Facility, protocol, and equipment

- (1) An SU will deliver most of its care to outpatients in addition to ED and inpatients
- (2) The SU should follow an internal protocol, which applies to diagnosis and management and is agreed by stakeholders
- (3) An equipped facility must be available
- (4) Essential equipment/tests:
 - 12-Lead ECG and 3-lead ECG monitoring
 - Non-invasive beat-to-beat blood pressure monitor with recording facilities for subsequent analysis
 - Tilt table
 - Holter monitors
 - External loop recorders
 - Follow-up of implantable loop recorders^a
 - 24-h blood pressure monitoring
 - Basic autonomic function tests
- (5) Established procedures for:
 - Echocardiography
 - Electrophysiological studies
 - Stress test
 - Neuroimaging tests
- (6) Specialists' consultancies (cardiology, neurology, internal medicine, geriatric, psychology), when needed

Therapy

Syncope patients will receive their therapy under the care of the SU unless expertise outside that of the unit is required.

Database management

The SU is required to keep medical records that should also include follow-up when appropriate. The database will also offer the possibility of collaborative research with other SUs.

^aImplantation of loop recorders may be performed either by SU physicians or by external cardiologists upon request of the SU physicians.

The staff

Most of the work is undertaken by nursing/technical staff. This requires specific skill and competence. In addition to assisting the syncope specialist, the specialized nurse/technician will perform procedures and tests (under physician supervision) provided that they are based on internal protocols and rules (Table 5).

Competence**Considerations**

Defining the area and level of competence of an SU is based on the following assumptions. Even if the skill of an individual syncope specialist may be insufficient to cover the whole case mix of the SU, the multidisciplinary skill of the different specialists involved in the SU should potentially be competent in all disorders referred to as T-LOC.

At present, the field lacks structured accreditation for clinical skills of the clinicians, and additional staff as well as equipment and facilities in the SU. The authors anticipate that this Position Paper will stimulate new structured training and accreditation opportunities. A diploma course, 'Syncope and Related disorders' for international participants, awarded by the Royal College of Physicians in Dublin, Ireland, is one example (www.rcpi.ie).

Disorders causing T-LOC are diverse and are reported to occur at different rates illustrating the scope of the necessary competencies. Results of 10 SUs^{13,17,19,23,41,45,55,56} (W. Wieling and J.G. van Dijk, personal communication) yielded the following categories (mean and range): reflex syncope (59%, 46–68), cardiac (10%, 1–35), orthostatic hypotension (9%, 1–19), unexplained syncope (11%, 5–18), psychogenic pseudosyncope (4%, 0–12), and epileptic seizures (1%, 0–5). Reflex syncope is by far the most common diagnosis. The rates of other disorders vary considerably, probably the result of differences in setting and specialty, and perhaps of limited diagnostic skills.

A second factor determining competence is risk management: the risks of cardiac syncope are high, those of reflex syncope low, with epilepsy, orthostatic hypotension and psychogenic T-LOC being intermediate. High risks require higher diagnostic skill levels regardless of frequency.

Thirdly, patient age affects the scope of competence. Syncope Units focusing on the elderly will encounter a different set of disorders compared with those with paediatric patients.⁴⁶ While paediatric T-LOC/syncope is not covered in depth by the 2009 ESC guidelines, the basic approach is the same as in adults, and children with T-LOC may profit as much from an SU as adults.

Syncope specialist

The considerations above prompted the following pragmatic description of a syncope specialist. A syncope specialist is a physician who has sufficient knowledge of historical clues and physical findings to recognize all major T-LOC forms, including mimics, as well as syndromes of orthostatic intolerance.

The 'syncope specialist'

The syncope specialist has responsibility for the comprehensive management of the patient from risk stratification to diagnosis, therapy, and follow-up, through a standardized protocol. The syncope specialist requires specific knowledge. The domains required are listed in the 'Competence' section.

Syncope specialists need not all have the same skill levels, but the SU as a whole must be able to provide a minimum skill set, so a combination of specialty skills is optimal. These conclusions are specified in *Consensus Statement 2—Competence and skills mix of physicians and staff required for syncope management in an SU*.

Notes on training

Syncope specialists typically start working in an SU after specialty training, so knowledge regarding T-LOC forms not covered by their specialty may need to be refreshed. Reflex syncope, orthostatic intolerance, and psychogenic pseudosyncope deserve special attention, as they usually do not routinely feature in prior training.

Reflex syncope is frequent in the population (30–40%) and in SUs (59%). Mastering the diagnosis of reflex syncope is difficult because its signs and symptoms are so variable that syncope cannot be defined practically using clinical descriptors: the ESC Guidelines defined T-LOC clinically but not syncope.⁵⁷ No ancillary test for reflex syncope meets the requirements,⁵⁸ leaving history taking as the prime diagnostic instrument. The importance of history taking and its high diagnostic yield⁴⁹ means that history taking should be allowed time. Syncope specialists typically set aside more time for history taking than novices, and may need up to 60 min to take a history and explain a diagnosis.^{58–62} As for risk prediction, rule sets did not perform better than clinical judgement.⁶³ A thorough knowledge of circulatory physiology helps to attribute historical elements to known circulatory patterns, strengthened by tilt table test experience.^{64,65} Psychogenic pseudosyncope occurs at vastly different frequencies in different studies. It can be recognized through history taking and often with tilt table testing.⁶⁶ Video-EEG monitoring is a preferable addition, conforming to the gold standard approach for psychogenic non-epileptic seizures.

Legend

The levels described here concern the SU as a whole, not those of individual physicians, except for column 1: the requested level of minimum basic diagnostic skills applies to each syncope specialist.

1. Diagnostic skills per syncope specialist

These skills refer to *history taking* and *physical examination* only.

'General medical knowledge': this is the level stipulated by medical education, i.e. the knowledge medical students are required to have in order to pass medical examinations; 'Specialist knowledge': this is the level required for specialty examination; when no specialty demands are available as for reflex syncope, the specialist level denotes the ability to recognize rare forms occurring once in about 200 cases.

2. SU access to ancillary tests

Ancillary diagnostic skills concern all necessary diagnostic tests.

Full access: the SU carries out the test.

Preferential access : the SU does not usually carry out the test but has preferential access to the tests/services in question.

3. SU ancillary tests

This is an overview of the most likely test required for the disorder in question.

Comprehensive care

The possibility of providing a comprehensive standardized syncope management of the patient from risk stratification to diagnosis, therapy, and follow-up characterizes and distinguishes an SU (*Figure 1*).


Referrals, access criteria

Referral can be directly from family practitioners, ED, in-hospital and out-hospital services, or self-referral from the patient. A fast-track access with separate waiting list and scheduled follow-up visits is recommended. In particular, patients at low/intermediate risk admitted to the ED should benefit from such a fast track

Consensus Statement 2—Competence and skills mix of physicians and staff required for syncope management in an SU

Major and minor category	1. Diagnostic skills per syncope specialist	2. SU access to ancillary tests	3. SU ancillary tests
Syncope			
Reflex	Specialist knowledge	Full	TT, CSM, 24 h BP, ELR-ILR
Syncope due to OH	Specialist knowledge	Full	TT, 24 h BP, autonomic testing
Cardiac syncope	Specialist knowledge	Full/preferential	ECG, telemetry, ELR-ILR, Echo, EPS
Epileptic seizures			
	General medical knowledge	Preferential	EEG, video-EEG monitoring, home video, neuroimaging
Psychogenic T-LOC			
PPS	General medical knowledge	Full/Preferential	TT, preferentially with video-EEG monitoring, home video
PNES	General medical knowledge	Preferential	video-EEG monitoring, home video

TT, tilt table test; CSM, carotid sinus massage; ILR, implantable loop recorder; ELR, external loop recorder; BP, blood pressure; Echo, echocardiogram; EEG, electroencephalogram; OH, orthostatic hypotension; PPS, psychogenic pseudosyncope; PNES, psychogenic non-epileptic seizure; EPS, electrophysiological testing.



(so-called 'protected discharge' or 'advanced access' with an appointment for early assessment), in order to reduce hospitalization rates, directly from the ED or after a short stay in the Short Observation unit of the ED as part of an internal protocol for risk stratification of intermediate risk ED patients.³⁶ In an existing multicentre experience,⁴¹ the majority of the patients (60%) were referred from out-of-hospital services, 11% had immediate referral and 13% delayed referral from the ED, and 16% were hospitalized. One of the critical variables for referrals from EDs is the initial clinical assessment by the non-specialist in syncope regarding the decision to admit. Since this issue is outside the scope of the present document, we refer to the existing guidelines, such as for example the ESC guidelines³ and the Canadian Specialist Consensus⁴ for defined algorithms on risk stratification and aligned actions regarding admission criteria.

Who should access the SU?

- Patients affected by T-LOC of suspected syncopal nature who, because of frequency or severity of the episodes, need to establish a diagnosis and to ascertain the need for a specific therapy.
- Patients with an already established diagnosis in order to receive a specialist's consultancy on the best evidence-based therapy or to start specific treatment, i.e. those listed in *Consensus Statement 4*.
- Patients who need follow-up to make a final diagnosis or assess efficacy of therapy (see 'Final report and follow-up' section)

Who should not access the SU?

- Patients with a certain diagnosis and/or an established indication for therapy, e.g. patients with bradycardia with guideline-based indications for cardiac pacing.
- Patients in whom syncope is a symptom secondary to underlying disease which requires urgent and specific diagnostic and therapeutic pathways that cannot be followed within an SU, e.g. syncope due to acute myocardial infarction or acute bleeding.

Only a minority of syncope patients seeking medical advice needs to be referred to an SU. In an existing multicentre experience,⁴¹ adult patients referred to an SU accounted for 163 per 100 000 inhabitants per year. The corresponding yearly incidence of syncope per 100 000 in the general population, ranges from 950 per 100 000 of the overall population of the Utah study⁶⁷ to 1100 per 100 000 in the patients >40 years old in the Framingham study.⁶⁸ By comparison, the yearly prevalence for patient visits to general practitioners is 930 per 100 000 inhabitants in a study performed in the Netherlands⁶ and 260 per 100 000 for patients referred to an ED in a multicentre Italian study.¹⁷

Management modalities

Diagnostic pathway

Diagnostic assessment consists of core assessment directly performed by the personnel of the SU and, when appropriate, by a fast-track access to any other tests and specialist visits that are not directly performed in the SU.

Consensus Statement 3—Tests and assessments available in SU

Initial assessment	History and physical evaluation including 3-min orthostatic BP measurement ^a 12-lead standard ECG
Subsequent tests and assessments (only when indicated)	
Blood tests	Electrolytes, haemoglobin, troponin, BNP, glucose, D-dimer, and haemogasanalysis/O ₂ saturation
Provocative tests	Carotid sinus massage and tilt table test
Monitoring	External loop recording, implantable loop recording, ambulatory 1–7 days ECG monitoring, and 24–48 h BP monitoring
Autonomic function tests	Standing test, Valsalva manoeuvre, deep breathing test, cold pressor test, and/or established procedures for access to other autonomic function tests
Cardiac evaluation	Established procedures for access to echocardiogram, stress test, electrophysiological study, and coronary angiography
Neurological evaluation	Established procedures for access to neurological tests (CT, MRI, EEG, video-EEG)
Geriatric evaluation	Established procedures for access to fall risk assessment (cognitive, gait and balance, visual, environmental) and for gait and balance retraining
Psychological or psychiatric evaluation	Established procedures for access to psychological or psychiatric consultancy (mental health problem or psychogenic syncope)

BNP, brain natriuretic peptide; BP, blood pressure; CT, computed tomography; ECG, electrocardiogram; EEG, electroencephalogram; MRI, magnetic resonance imaging; SU, Syncope Unit.
^aPostural orthostatic tachycardia may require longer stands.

Treatment pathway

Even if the model of therapy delivery is that which is most appropriate to existing practice and resources, those listed in *Consensus Statement 4* are the minimum core treatments that should be provided directly by the SU staff.

Consensus Statement 4—Treatments available in SU

Reflex syncope	<ul style="list-style-type: none"> Structured education on management of risk factor for prevention of reflex syncope (including adequate fluid and salt intake) Biofeedback training^a and instruction sheet on counter pressure manoeuvres and leg/abdomen elastic stocking Established procedures for access to pacemaker implantation Modification of culprit drugs Prescription of vasoconstrictor drugs
Orthostatic hypotension	<ul style="list-style-type: none"> Structured education session on management of risk factor for prevention of orthostatic hypotension (including adequate fluid and salt intake and care during orthostatic changes after prolonged periods lying or sitting) Leg/abdomen elastic stocking Modification of culprit hypotensive drugs Prescription of antihypotensive drugs, vasoconstrictor drugs

Continued

Cardiac syncope	<ul style="list-style-type: none"> Established procedures for access to pacemaker implantation, catheter ablation of arrhythmias, ICD therapy and other surgical interventions Prescription of antiarrhythmic drugs
Falls	Structured education on management of risk factor for fall prevention
Psychogenic pseudosyncope (functional)	Established procedures for access to psychotherapy
Epilepsy	Established procedures for access to neurology care

ICD, Implantable cardioverter-defibrillator.
^aBiofeedback means that the training session of the counter pressure manoeuvres consists of biofeedback training using a continuous blood pressure monitor. Each manoeuvre is demonstrated and explained. The manoeuvres are practiced under supervision, with immediate feedback of the recordings to gain optimal performance.⁵⁴

Final report and follow-up

Once the work-up is complete and a therapy is prescribed, the patient and referring practitioner should receive a written comprehensive final report and be referred back to the referring physician. The framework of a structured final report is described in *Tables 6 and 7*.

Table 6 Framework of a structured final report (clinic note)

Chief complaint	Use definition recommended by the ESC guidelines on syncope reported in <i>Tables 7</i>
Present illness	<p>Circumstances just prior the T-LOC event (predisposing factors, triggers, body position, activity, signs and symptoms of autonomic activation, accompanying symptoms, prodromes)</p> <p>Features of the attack (eyewitness), duration, recovery, consequences (trauma)</p> <p>Previous T-LOC episodes (number, date of occurrence, features, previous tests and treatments)</p>
Past medical history	<p>Description of the underlying diseases</p> <p>Medications</p> <p>Social history</p> <p>Family history</p>
Physical examination	Findings
Electrocardiogram	Findings
Tests	List and report of the relevant findings of the tests performed
Assessment and diagnosis	Description of the findings that led to the final diagnosis
Plan/treatment	Suggested further diagnostic tests or treatment with an explanation of the expected results, as appropriate
Follow-up modality	Description and eventual appointment for further visits

Table 7 Classification and criteria used to establish a certain/likely diagnosis (derived from the 2009 ESC guidelines) on syncope to be reported in the final report

Classification of T-LOCs	Diagnostic features
Reflex syncope (neurally mediated)	
Classical Vasovagal Syncope	If the syncope was precipitated by emotional distress (fear, severe pain, medical instrumentation procedures) or prolonged standing and was associated with typical prodromes
Carotid sinus syncope	If the syncope was reproduced by 10 s of bilateral sequential carotid sinus massage, supine, and erect, in the presence of asystole > 3 s and/or fall in systolic blood pressure > 50 mmHg and in the absence of competing diagnoses
Situational syncope	If the syncope occurred during or immediately after urination, defecation, coughing, laughing, or swallowing, etc.
Atypical form	If the syncope occurred with uncertain or even apparently absent triggers. The diagnosis then rests less on history taking alone, and more on the exclusion of other causes of syncope (the absence of structural heart disease) and on reproducing similar symptoms with tilt testing
Orthostatic intolerance	
Orthostatic hypotension	If syncope occurred after standing up and symptomatic orthostatic hypotension was documented. In the 'Initial form', orthostatic hypotension occurs immediately on standing then spontaneously and rapidly returns to normal within 30 s. In the <i>Classical form</i> orthostatic hypotension occurs after 30 s but within 3 min after active standing up, while in <i>Progressive (delayed) form</i> —usually diagnosed by means of tilt testing—progressive orthostatic hypotension occurs more than 3 min after standing up
Postural tachycardia syndrome (POTS)	If the symptoms of orthostatic intolerance are precipitated by an increase in heart rate > 30 b.p.m. (> 40 b.p.m. in patients 12–19 years old) vs. supine and the absence of orthostatic hypotension
Cardiac	
Cardiac arrhythmia	If the class I diagnostic criteria of the ESC guidelines ^{1,2} were met during prolonged ECG monitoring or by means of electrophysiological study; cardiac arrhythmia also included the case of patients with severely depressed EF who had a definite indication for ICD regardless of the mechanism of syncope
Structural cardiac or cardiopulmonary	If the patient was affected by acute cardiac ischaemia or other acute cardiopulmonary diseases or prolapsing atrial myxoma or severe aortic stenosis or hypertrophic obstructive cardiomyopathy, etc.
Non-syncopal attacks	
Epilepsy, unexplained fall, psychogenic pseudosyncope (functional), vertebro-basilar, metabolic disorders, etc.	If the episode of T-LOC was initially attributed to a syncopal condition but the subsequent evaluation demonstrated a non-syncopal mechanism (i.e. epilepsy, vertebro-basilar ischaemic attack, unexplained fall, or psychogenic [functional] pseudosyncope), hypoxia, intoxication, cataplexy, drop attacks)
Unexplained syncope/T-LOC	In those patients without any of the above diagnosis

However, the patient should still be periodically followed-up by the SU staff for three main purposes:

1. to assess the efficacy of treatment and re-evaluation in the case of syncope recurrence,
2. to establish a diagnosis in those patients with unexplained syncope who are undergoing prolonged ECG monitoring with external or implantable loop recorders, and
3. to contribute to the program of quality control of the SU (follow-up database, audit, etc.).

The follow-up process can be easily performed via phone interview or other tools, e.g. web-based.

Quality indicators

Based on the data presented in the previous sections, this Task Force believes that reasonable goals (expected benefits) of the EHRA SU should be those listed in the *Consensus Statement 5*. Since clinical outcomes will vary according to the local patient

cohorts in different geographic locations, the method of comparative effectiveness to assess the relevant clinical outcomes before and after the implementation of the SU has been considered. The minimum acceptable values should be compared with usual care without SU. Published information to serve as baseline data may be used if local information is not available (e.g. national data, or hospital practice before establishing the SU, or practice with similar characteristics in nearby hospitals without SU). As the practice of the SU evolves, new outcome metrics will be developed; the below suggested metrics are recommendation for a starting SU. Alternative approaches may include reduction in unnecessary/inappropriate investigations; the ability of individual practices to achieve these targets will depend on data available and local practices. Whereas the guidelines are aiming to standardize syncope management, this TF acknowledge that it may take some years to achieve this depending on maturity of services and local practice.

The objectives of the suggested metrics are: (i) to facilitate cost-effectiveness, (ii) to prepare future accreditation for SUs and training for staff, and (iii) to establish benchmark for quality control. However,

the metrics are not intended to be imperative for practice but rather a rough guide and dependent on extant practice, adequate human and other resources and the specific healthcare setting. These are advisory metrics and should not be adopted as mandatory targets until further research on preferred syncope management structures is conducted relevant to different countries and healthcare systems. Furthermore, the current field lacks structured

accreditation programmes for T-LOC/Syncope and, until these are established and widely available, specified training metrics are difficult to implement. The details of these metrics should be reviewed and approved by the lead physician, the syncope team, the hospital clinical practice committee, and stakeholders. External or internal audit via a registry mechanism may be considered to evaluate the quality outcomes.

Consensus Statement 5—Quality indicators

Quality indicator	Process indicator	Desirable outcome target	Comments
1. SU			
To reduce the rate of unexplained T-LOC	At least 70% of patients receive a definite diagnosis (according to ESC guidelines definitions)	Absolute rate of unexplained T-LOC $\leq 20\%$	Data from published literature indicate that a reasonable goal is to achieve an absolute rate of unexplained T-LOC $\leq 20\%$ (see 'Rationale for an SU' section)
To reduce the rate of hospitalization (in patients at intermediate–high risk from ED)	At least 20% of patients with unexplained syncope after initial ED evaluation have fast-track access to SU for early assessment	$\leq 20\%$ of patients with unexplained T-LOC admitted after ED initial evaluation (according to ESC guidelines definition)	Data from published literature indicate that a reasonable goal is to achieve an absolute rate of admissions $\leq 20\%$ (see 'Rationale for an SU' section)
To reduce costs per patient	At least 20% reduction in costs relative to usual local practice		Applies to new services
To improve the outcome	Less than 5% re-admissions for syncope recurrence in patients with an established and successfully treated diagnosis (according to ESC guidelines definitions)	Less than 20% of paced patients have recurrence of syncope at 1 year	Older patients who have more than one attributable cause of T-LOC may experience recurrence of symptoms due to other causes such as autonomic dysfunction and medication
2. Personnel			
Leadership	One physician leading the SU		The leader is usually one of the Syncope Specialists of the SU (see Specialist Consensus Statement 1).
Staff	A dedicated trained staff (specifics for training should map the T-LOC diagnoses and treatments)		The credentialing and the number of staff would vary according to the size and organization of the local practice
Training	Syncope Specialist and Staff attend regular training programme and conferences in addition to accreditation programme		At present, the field lacks structured accreditation for T-LOC/Syncope
3. Operations			
Number of patients	At least 100 new cases per year per SU		This is the minimum number necessary in order to develop and maintain expertise for one syncope specialist and one staff personnel and for the SU to be cost-efficient
Tests	>95% of patients have a documented ECG >90% of patients have documented orthostatic tests >90% have carotid sinus massage, tilt table test, external loop recorder and implantable loop recorder performed according to ESC guidelines indications		
Waiting list (first visit and follow-up)	70% of low risk patients seen within 3 months 90% of intermediate risk patients seen within 2 weeks No waiting list for high-risk patients		

EHRA Scientific Document Committee, Gregory Y. H. Lip (EHRA Scientific Documents Committee Chair), Bulent Gorenek (EHRA Scientific Documents Committee Co-Chair), Christian Sticherling, Laurent Fauchier, Hein Heidbuchel, Angel Moya Mitjans, Marc A. Vos, Michele Brignole, Gheorghe-Andrei Dan, Michele Gulizia, Francisco Marin, Giuseppe Boriani, Deirdre Lane, and Irene Savelieva.

References

- Brignole M, Alboni P, Benditt D, Bergfeldt L, Blanc JJ, Bloch Thomsen PE *et al*. Task Force on Syncope, European Society of Cardiology. Guidelines on management (diagnosis and treatment) of syncope. *Eur Heart J* 2001;**22**:1256–306.
- Brignole M, Alboni P, Benditt DG, Bergfeldt L, Blanc JJ, Bloch Thomsen PE *et al*. Guidelines on management (diagnosis and treatment) of syncope—update 2004. *Eurpace* 2004;**6**:467–537.
- Moya A, Sutton R, Ammirati F, Blanc JJ, Brignole M, Dahm JB *et al*. Guidelines for the diagnosis and management of syncope (version 2009). *Eur Heart J* 2009;**30**:2631–71.
- Sheldon RS, Morillo CA, Krahn AD, O'Neill B, Thiruganasambandamoorthy V, Parkash R *et al*. Standardized approaches to the investigation of syncope: Canadian Cardiovascular Society position paper. *Can J Cardiol* 2011;**27**:246–53.
- Brignole M, Hamdan MH. New concepts in the assessment of syncope. *J Am Coll Cardiol* 2012;**59**:1583–91.
- Olde Nordkamp LR, van Dijk N, Ganzeboom KS, Reitsma JB, Luitse JS, Dekker LR *et al*. Syncope prevalence in the ED compared to general practice and population: a strong selection process. *Am J Emerg Med* 2009;**27**:271–9.
- Casini-Raggi V, Bandinelli G, Lagi A. Vasovagal syncope in emergency room patients: analysis of a metropolitan area registry. *Neuroepidemiology* 2002;**21**:287–91.
- Kenny RA, Bhangu J, King-Kallimanis BL. Epidemiology of syncope/collapse in younger and older Western patient populations. *Prog Cardiovasc Dis* 2013;**55**:357–63.
- Baron-Esquivias G, Moreno SG, Martinez A, Pedrote A, Vazquez F, Granados C *et al*. Cost of diagnosis and treatment of syncope in patients admitted to a cardiology unit. *Eurpace* 2006;**8**:122–7.
- Krahn AD, Klein GJ, Yee R, Hoch JS, Skanes AC. Cost implications of testing strategy in patients with syncope - randomized assessment of syncope trial. *J Am Coll Cardiol* 2003;**42**:495–501.
- Kenny RA, Rice C, Byrne L. The role of the Syncope Management Unit. *Card Electrophysiol Clin* 5:529–42.
- Kenny RA, O'Shea D, Walker HF. Impact of a dedicated syncope and falls facility for older adults on emergency beds. *Age Ageing* 2002;**31**:272–5.
- Brignole M, Disertori M, Menozzi C, Raviele A, Alboni P, Pitzalis MV *et al*. Management of syncope referred urgently to general hospitals with and without syncope units. *Eurpace* 2003;**5**:293–8.
- Farwell DJ, Sulke AN. Does the use of a syncope diagnostic protocol improve the investigation and management of syncope? *Heart* 2004;**90**:52–8.
- Shen WK, Decker WW, Smars PA, Goyal DG, Walker AE, Hodge DO *et al*. Syncope Evaluation in the Emergency Department Study (SEEDS): a multidisciplinary approach to syncope management. *Circulation* 2004;**110**:3636–45.
- Blanc JJ, L'Her C, Gosselin G, Cornily JC, Fatemi M. Prospective evaluation of an educational programme for physicians involved in the management of syncope. *Eurpace* 2005;**7**:400–6.
- Brignole M, Ungar A, Bartoletti A, Ponassi I, Lagi A, Mussi C *et al*. Standardized-care pathway vs. usual management of syncope patients presenting as emergencies at general hospitals. *Eurpace* 2006;**8**:644–50.
- Parry SW, Reeve P, Lawson J, Shaw FE, Davison J, Norton M *et al*. The Newcastle protocols 2008: an update on head-up tilt table testing and the management of vasovagal syncope and related disorders. *Heart* 2009;**95**:416–20.
- Ammirati F, Colaceci R, Cesario A, Strano S, Della Scala A, Colangelo I *et al*. Management of syncope: clinical and economic impact of a Syncope Unit. *Eurpace* 2008;**10**:471–6.
- Fedorowski A, Burri P, Juul-Moller S, Melander O. A dedicated investigation unit improves management of syncopal attacks (Syncope Study of Unselected Population in Malmo--SYSTEMA I). *Eurpace* 2010;**12**:1322–8.
- McCarthy F, De Bhaldraithe S, Rice C, McMahon CG, Geary U, Plunkett PK *et al*. Resource utilisation for syncope presenting to an acute hospital Emergency Department. *Ir J Med Sci* 2010;**179**:551–5.
- Daccarett M, Jetter TL, Wasmund SL, Brignole M, Hamdan MH. Syncope in the emergency department: comparison of standardized admission criteria with clinical practice. *Eurpace* 2011;**13**:1632–8.
- Shin TG, Kim JS, Song HG, Jo JJ, Sim MS, Park SJ. Standardized approaches to syncope evaluation for reducing hospital admissions and costs in overcrowded emergency departments. *Yonsei Med J* 2013;**54**:1110–8.
- Sun BC. Quality-of-life, health service use, and costs associated with syncope. *Prog Cardiovasc Dis* 2013;**55**:370–5.
- Sanders NA, Jetter TL, Brignole M, Hamdan MH. Standardized care pathway versus conventional approach in the management of patients presenting with faint at the University of Utah. *Pacing Clin Electrophysiol* 2013;**36**:152–62.
- Sun BC, McCreath H, Liang LJ, Bohan S, Baugh C, Ragsdale L *et al*. Randomized clinical trial of an emergency department observation syncope protocol versus routine inpatient admission. *Ann Emerg Med* 2014;**64**:167–75.
- Rauci U, Scateni S, Tozzi AE, Drago F, Giordano U, Marcias M *et al*. The availability and the adherence to pediatric guidelines for the management of syncope in the Emergency Department. *J Pediatr* 2014;**165**:967–72 e1.
- Josephson CB, Rahey S, Sadler RM. Neurocardiogenic syncope: frequency and consequences of its misdiagnosis as epilepsy. *Can J Neural Sci* 2007;**34**:221–4.
- Zaidi A, Clough P, Cooper P, Scheepers B, Fitzpatrick AP. Misdiagnosis of epilepsy: many seizure-like attacks have a cardiovascular cause. *J Am Coll Cardiol* 2000;**36**:181–4.
- Passman R, Horvath G, Thomas J, Kruse J, Shah A, Goldberger J *et al*. Clinical spectrum and prevalence of neurologic events provoked by tilt table testing. *Arch Intern Med* 2003;**163**:1945–8.
- MacCormick JM, McAlister H, Crawford J, French JK, Crozier I, Shelling AN *et al*. Misdiagnosis of long QT syndrome as epilepsy at first presentation. *Ann Emerg Med* 2009;**54**:26–32.
- Bartoletti A, Fabiani P, Gianni R, Cappelletti C, Santoro GM, Fortini A *et al*. After the 2001 European Society of Cardiology Guidelines: is it possible to reduce the number of patients with syncope to be hospitalized?. *Minerva Med* 2004;**95**:451–60.
- Iyer PM, McNamara PH, Fitzgerald M, Smyth L, Dardis C, Jawad T *et al*. A seizure care pathway in the emergency department: preliminary quality and safety improvements. *Epilepsy Res Treat* 2012;**2012**:273175.
- Sun BC, Emond JA, Camargo CA Jr. Direct medical costs of syncope-related hospitalizations in the United States. *Am J Cardiol* 2005;**95**:668–71.
- Parry SW, Frearson R, Steen N, Newton JL, Tryambake P, Kenny RA. Evidence-based algorithms and the management of falls and syncope presenting to acute medical services. *Clin Med* 2008;**8**:157–62.
- Ungar A, Tesi F, Chisciotti VM, pepe G, Vanni S, Grifoni S *et al*. Assessment of a novel management pathway for patients referred to the Emergency Department for syncope: results in a tertiary hospital. *Eurpace* 2015;doi:10.1093/europace/euv106.
- Calkins H, Byrne M, el-Atassi R, Kalbfleisch S, Langberg JJ, Morady F. The economic burden of unrecognized vasodepressor syncope. *Am J Med* 1993;**95**:473–9.
- Steinberg LA, Knilans TK. Costs and utility of tests in the evaluation of the pediatric patients with syncope. *Prog Pediatr Cardiol* 2001;**13**:139–49.
- Disertori M, Brignole M, Menozzi C, Raviele A, Rizzon P, Santini M *et al*. Management of patients with syncope referred urgently to general hospitals. *Eurpace* 2003;**5**:283–91.
- Edvardsson N, Frykman V, van Mechelen R, Mitro P, Mohii-Oskarsson A, Pasquie JL *et al*. Use of an implantable loop recorder to increase the diagnostic yield in unexplained syncope: results from the PICTURE registry. *Eurpace* 2011;**13**:262–9.
- Brignole M, Ungar A, Casagrandi I, Gulizia M, Lunati M, Ammirati F *et al*. Prospective multicentre systematic guideline-based management of patients referred to the Syncope Units of general hospitals. *Eurpace* 2010;**12**:109–18.
- Brignole M, Malasana G, Sherwood RP, Daccarett M, Jetter TL, Hamdan MH. Evaluation of patients with "faint" in an American teaching hospital: a dire need for a standardized approach. *Pacing Clin Electrophysiol* 2011;**34**:284–90.
- Croci F, Brignole M, Alboni P, Menozzi C, Raviele A, Del Rosso A *et al*. The application of a standardized strategy of evaluation in patients with syncope referred to three syncope units. *Eurpace* 2002;**4**:351–5.
- Ungar A, Del Rosso A, Giada F, Bartoletti A, Furlan R, Quartieri F *et al*. Early and late outcome of treated patients referred for syncope to emergency department: the EGSYS 2 follow-up study. *Eur Heart J* 2010;**31**:2021–6.
- Sousa, Marques, Faria. Syncope unit: Experience of a center using diagnostic flow-charts for syncope of uncertain etiology after initial assessment. *Rev Port Cardiol* 2012;**32**:581–91.
- Newton JL, Marsh A, Frith J, Parry S. Experience of a rapid access blackout service for older people. *Age Ageing* 2010;**39**:265–8.
- Fitzpatrick AP, Cooper P. Diagnosis and management of patients with blackouts. *Heart* 2006;**92**:559–68.
- Petkar S, Bell W, Rice N, Iddon P, Cooper P, McKee D *et al*. Initial experience with a rapid access blackouts triage clinic. *Clin Med* 2011;**11**:11–6.
- van Dijk N, Boer KR, Colman N, Bakker A, Stam J, van Grieken JJ *et al*. High diagnostic yield and accuracy of history, physical examination, and ECG in patients with transient loss of consciousness in FAST: the Fainting Assessment study. *J Cardiovasc Electrophysiol* 2008;**19**:48–55.
- Vieling W, van Dijk N, de Lange FJ, Olde Nordkamp LR, Thijs RD, van Dijk JG *et al*. History taking as a diagnostic test in patients with syncope: developing expertise in syncope. *Eur Heart J* 2015;**36**:277–80.
- Rodriguez-Entem F, Gonzalez-Enriquez S, Olalla-Antolin JJ, Cobo-Belaustegui M, Exposito-Garcia V, Llano-Cardenal M *et al*. [Management of syncope in the

- emergency department without hospital admission: usefulness of an arrhythmia unit coordinated protocol]. *Rev Esp Cardiol* 2008;**61**:22–8.
52. Brignole M, Menozzi C, Bartoletti A, Giada F, Lagi A, Ungar A et al. A new management of syncope: prospective systematic guideline-based evaluation of patients referred urgently to general hospitals. *Eur Heart J* 2006;**27**:76–82.
 53. Wieling W, van Dijk N, de Lange FJ, Olde Nordkamp LR, Thijs RD, van Dijk JG et al. History taking as a diagnostic test in patients with syncope: developing expertise in syncope. *Eur Heart J* 2015;**36**:277–80.
 54. van Dijk N, Quartieri F, Blanc JJ, Garcia-Civera R, Brignole M, Moya A et al. Effectiveness of physical counterpressure maneuvers in preventing vasovagal syncope: the Physical Counterpressure Manoeuvres Trial (PC-Trial). *J Am Coll Cardiol* 2006;**48**:1652–7.
 55. Strano S, Colosimo C, Sparagna A, Mazzei A, Fattouch J, Giallonardo AT et al. Multidisciplinary approach for diagnosing syncope: a retrospective study on 521 outpatients. *J Neurol Neurosurg Psychiatry* 2005;**76**:1597–600.
 56. Mitro P, Kirsch P, Valocik G, Murin P. A prospective study of the standardized diagnostic evaluation of syncope. *Europace* 2011;**13**:566–71.
 57. Saal DP, van Dijk JG. Classifying syncope. *Auton Neurosci* 2014;**184**:3–9.
 58. Sheldon R. Tilt table testing and implantable loop recorders for syncope. *Cardiol Clin* 2013;**31**:67–74.
 59. Sheldon R, Rose S, Connolly S, Ritchie D, Koshman ML, Frenneaux M. Diagnostic criteria for vasovagal syncope based on a quantitative history. *Eur Heart J* 2006;**27**:344–50.
 60. Sheldon R. Syncope diagnostic scores. *Prog Cardiovasc Dis* 2013;**55**:390–5.
 61. Thijs RD, Bloem BR, van Dijk JG. Falls, faints, fits and funny turns. *J Neurol* 2009;**256**:155–67.
 62. Brignole M, Vardas P, Hoffman E, Huikuri H, Moya A, Ricci R et al. Indications for the use of diagnostic implantable and external ECG loop recorders. *Europace* 2009;**11**:671–87.
 63. Costantino G, Casazza G, Reed M, Bossi I, Sun B, Del Rosso A et al. Syncope risk stratification tools vs clinical judgment: an individual patient data meta-analysis. *Am J Med* 2014;**127**:1126 e13–25.
 64. Wieling W, Thijs RD, van Dijk N, Wilde AA, Benditt DG, van Dijk JG. Symptoms and signs of syncope: a review of the link between physiology and clinical clues. *Brain* 2009;**132**(Pt 10):2630–42.
 65. van Dijk JG, Thijs RD, van Zwet E, Tannemaat MR, van Niekerk J, Benditt DG et al. The semiology of tilt-induced reflex syncope in relation to electroencephalographic changes. *Brain* 2014;**137**(Pt 2):576–85.
 66. Tannemaat MR, van Niekerk J, Reijntjes RH, Thijs RD, Sutton R, van Dijk JG. The semiology of tilt-induced psychogenic pseudosyncope. *Neurology* 2013;**81**:752–8.
 67. Malasana G, Brignole M, Daccarett M, Sherwood R, Hamdan MH. The prevalence and cost of the faint and fall problem in the state of Utah. *Pacing Clin Electrophysiol* 2011;**34**:278–83.
 68. Soteriades ES, Evans JC, Larson MG, Chen MH, Chen L, Benjamin EJ et al. Incidence and prognosis of syncope. *N Engl J Med* 2002;**347**:878–85.

Appendix

Development process of the document

The Task Force members constituted the writing group aimed to provide a succinct, evidence-based document at a uniform level, rather than a comprehensive narrative review. The Task Force members met for the first time on October 2013; they prepared the table of contents and assigned each section to compact writing groups with two members who completed the first versions and developed preliminary consensus statements. A second plenary meeting, which was open also to the External Contributors, was held in June 2014 during which the consensus statements were agreed and voted upon. The provisional draft of the entire document was then submitted to External Contributors who provided written comments. To establish consensus, we conducted surveys of the entire group (writing members and external contributors) using a pre-defined threshold for agreement as a vote of >80% for each Consensus Statement.

The specific wording of definitions, statements, and the choice of references were the result of prolonged debate and consensus-seeking. The text underwent iterative revisions to resolve differences, increase clarity, and align the document format. The chair and co-chair took the responsibility to manage the process and wrote the final version of the text that was approved by the members of the Task Force and by the External Contributors.